

AD-A114 326

TEXAS UNIV AT AUSTIN CENTER FOR CYBERNETIC STUDIES F/8 12/1
KHINCIN-KULLBACK-LEIBLER ESTIMATION WITH INEQUALITY CONSTRAINTS-ETC(U)
NOV 81 A CHARNES, W W COOPER, J TYSSDAL N00014-75-C-0569
CCS-414 NL

UNCLASSIFIED

For
AD A
7 5800



END
DATE
FILMED
5-82
DTIC

PA 11/4

CENTER FOR CYBERNETIC STUDIES

The University of Texas
Austin, Texas 78712

DTIC
ELECTE
MAY 5 1982
S H

STATEMENT A
public release
Unlimited

82 05 04

Research Report CCS 414

12

^Y
KHINČIN-KULLBACK-LEIBLER ESTIMATION
WITH INEQUALITY CONSTRAINTS

by

A. Charnes
W.W. Cooper
J. Tyssedal*

November 1981

*University of Bergen, Norway

This research was partly supported by ONR Contract N00014-75-C-0569 with the Center for Cybernetic Studies, The University of Texas at Austin. Reproduction in whole or in part is permitted for any purpose of the United States Government.

CENTER FOR CYBERNETIC STUDIES

A. Charnes, Director
Business-Economics Building, 203E
The University of Texas at Austin
Austin, Texas 78712
(512) 471-1821

DISTRIBUTION STATEMENT A
Approved for public release
Distribution Unlimited

DTIC
COLLECTED
MAY 5 1982
H

ABSTRACT

In this paper we extend our grasp of statistical theory, duality theory, and computational convenience to the general linear inequality situation in K^2L (or minimum discrimination information) estimation by exhibiting it as the limit of a simple one parameter sequence of equality problems. Only the finite discrete distribution case is treated here.

KEY WORDS

Khinchin-Kullback-Leibler Estimation

Inferential Distribution

Minimum Discrimination Information

Inequality Constrained Distributions



Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
<input type="checkbox"/>	<input type="checkbox"/>

INTRODUCTION

Frequently in hypothesis testing or estimation of an "inferential" distribution (in Akaike's terminology) by the Khinčin-Kullback-Leibler (or Minimum Discrimination Information) method one has information about the possible candidate distributions in the form of linear inequalities on the components of the distribution in addition to equality (moment) constraints. In MOFS 1978, Charnes, Cooper, and Seiford [1] developed a convex programming duality theory for the K^2L method with linear inequality constraints. It is especially incisive and convenient for constraints in equality (or moment) form which, further, have connections with established statistical theory as well as the analytic and computational facility of an unconstrained extremal problem in simple smooth functions.

In this paper we extend our grasp of statistical theory, duality theory, and computational convenience to the general linear inequality situation by exhibiting it as the limit of a simple one parameter sequence of equality problems. We treat only the finite discrete distribution case here, reserving the general distribution case to a forthcoming paper which also simplifies the duality theory of the "equality" case for general distributions as developed by Ben-Tal and Charnes in [2].

INEQUALITY AND APPROXIMATING EQUALITY FORMS

The dual programs for the inequality form as presented in Charnes-Cooper-Seiford [1] are

$$\begin{aligned} \max \quad v(\delta) &\triangleq -\delta^T \ln \left[\frac{\delta}{ec} \right] \\ \text{(I)} \quad \text{s.t.} \quad &\delta^T A^1 = b^{1T} \\ &\delta^T A^2 + \gamma^T = b^{2T} \end{aligned}$$

$$\delta, \gamma \geq 0$$

and

$$\begin{aligned} \text{(II)} \quad \inf \xi(z) &\triangleq c^T e A^1 z^1 + A^2 z^2 - b^{1T} z^1 - b^{2T} z^2 \\ \text{s.t.} \quad &-z^2 \geq 0 \end{aligned}$$

To obtain the dual programs for our approximating (weighted) equality form, we employ the procedure in [1]. Thus, consider

$$(1) \quad K(x, y, \delta, \gamma) \triangleq \sum_{i \in I^1} (c_i e^{x_i} - \delta_i x_i) + \sum_{i \in I^2} (\epsilon c_i e^{(y_i/\epsilon)} - \gamma_i y_i)$$

where $c_i > 0$ for all i and $\epsilon > 0$.

Minimizing with respect to x_i and y_i gives:

$$(2) \quad x_i^* = \ln \left(\frac{\delta_i}{c_i} \right), \quad i \in I^1, \quad y_i^* = \epsilon \ln \left(\frac{\gamma_i}{\epsilon c_i} \right), \quad i \in I^2$$

Hence,

$$(3) \quad -\sum_{i \in I^1} \delta_i \ln \left(\frac{\delta_i}{c_i} \right) - \epsilon \sum_{i \in I^2} \gamma_i \ln \left(\frac{\gamma_i}{\epsilon c_i} \right) \leq K(x, y, \delta, \gamma)$$

To decouple, as in [1], we obtain

$$(4) \quad \delta^T x + \gamma^T y = b^{1T} z^1 + b^{2T} z^2$$

and

$$(5) \quad (\delta^T, \gamma^T) \begin{bmatrix} A^1 & A^2 \\ 0 & I \end{bmatrix} \begin{bmatrix} z^1 \\ z^2 \end{bmatrix} = (b^{1T}, b^{2T}) \begin{bmatrix} z^1 \\ z^2 \end{bmatrix}$$

when we choose

$$(6) \quad x = A^1 z^1 + A^2 z^2, \quad y = z^2$$

Thereby, we have the dual problems

$$(III) \quad \max_{\delta, \gamma, \varepsilon} v(\delta, \gamma, \varepsilon) \triangleq - \sum_{i \in I^1} \delta_i \ln \left(\frac{\delta_i}{\varepsilon c_i} \right) - \varepsilon \sum_{i \in I^2} \gamma_i \ln \left(\frac{\gamma_i}{\varepsilon c_i} \right)$$

s.t.

$$\delta^T A^1 = b^1 T$$

$$\delta^T A^2 + \gamma^T = b^2 T$$

$$\delta, \gamma \geq 0$$

and

$$(IV) \quad \inf \xi(z) \triangleq c^1 T_e A^1 z^1 + A^2 z^2 - b^1 T_z^1 - b^2 T_z^2 + \varepsilon c^2 T_e (z^2 / \varepsilon)$$

with z unconstrained

(Note that the $c_i, i \in I_2$, may be chosen arbitrarily.)

The duality theory of (III) and (IV) is precisely that of the equality case in [1], as may be seen by making the change of variables to

$$\tilde{\gamma}_1 \triangleq \varepsilon \gamma_1, \quad \tilde{c}_1 \triangleq \begin{bmatrix} c_i, i \in I^1 \\ \varepsilon c_i, i \in I^2 \end{bmatrix}.$$

Our present form is, however, more convenient for our arguments.

We now define

$$(7.1) \quad f(z) \triangleq c^1 T_e A^1 z^1 + A^2 z^2 - b^1 T_z^1 - b^2 T_z^2, \quad z^2 \leq 0$$

$$(7.2) \quad g(z, \varepsilon) \triangleq c^1 T_e A^1 z^1 + A^2 z^2 - b^1 T_z^1 - b^2 T_z^2 + \varepsilon c^2 T_e (z^2 / \varepsilon)$$

We will then have

Theorem 1:

For some $\epsilon > 0$, $\inf_{z^2 \leq 0} f(z)$ exists if and only if $\inf_z g(z, \epsilon)$ exists.

Further,

$$\lim_{\epsilon \rightarrow 0} \inf_z g(z, \epsilon) = \inf_{z^2 \leq 0} f(z)$$

Proof:

Consider $\frac{\partial g}{\partial z_i^2}$. It may be written as

$$(8) \quad \frac{\partial g}{\partial z_i^2} = (c^1_k)^T e^{A^1 z^1 + A^2 z^2 - b^2_1 + z^2_1 c_1 e^{(z^2_1/\epsilon)}} \quad , \quad i \in I^2$$

where $k_r \triangleq \frac{\partial}{\partial z_i^2} (r A^2 z^2)$ and $(c^1_k)^T \triangleq (c^1_{k_1}, \dots, c^1_{k_m})$.

For $z^2_1 > 0$ we see that by choosing ϵ small enough we can make $g(z, \epsilon)$ an increasing function in that direction. Thus, in seeking a minimum we need only consider $z^2 \leq 0$. For notational simplicity in the following and "inf" with respect to z shall always also entail $z^2 \leq 0$.

For $z^2 \leq 0$ and all $\epsilon > 0$ we have

$$(9) \quad f(z) \leq g(z, \epsilon) \leq f(z) + \epsilon \sum_{i \in I^2} c_i$$

If $\inf f(z)$ exists, then, by (9), $\inf f(z) \leq g(z, \epsilon)$ so that $\inf g(z, \epsilon)$ exists for all $\epsilon > 0$. Further, $\inf f(z) \leq \inf g(z, \epsilon)$. On the other hand, if $\inf g(z, \epsilon)$ exists, then, by (9), $\inf g(z, \epsilon) \leq f(z) + \epsilon \sum_{i \in I^2} c_i$ so that $\inf f(z)$ exists. Further, too, $\inf g(z, \epsilon) \leq \inf f(z) + \epsilon \sum_{i \in I^2} c_i$.

Hence

$$(10.1) \quad \inf f(z) \leq \inf g(z, \varepsilon) \leq \inf f(z) + \varepsilon \sum_{i \in I^2} c_i$$

and

$$(10.2) \quad \lim_{\varepsilon \rightarrow 0} \inf g(z, \varepsilon) = \inf f(z)$$

When $f(z)$ has a minimum at z^* , again from (9) we can conclude

$$(11) \quad f(z^*) = \lim_{\varepsilon \rightarrow 0} g(z^*, \varepsilon)$$

Q.E.D.

So much for the minimization side of the duality. For the K^2_L side we have

Theorem 2:

The maximum in problem (I) and, as $\varepsilon \rightarrow 0$, in problem (III) is the same.

Proof:

For $0 \leq \gamma_i \leq \varepsilon c_i$, we have $-c_i \leq \gamma_i \ln\left(\frac{\gamma_i}{\varepsilon c_i}\right) \leq 0$, whereas for

$\gamma_i > \varepsilon c_i$, $\gamma_i \ln\left(\frac{\gamma_i}{\varepsilon c_i}\right) > 0$. Therefore,

$$(12) \quad -\varepsilon \sum_{i \in I^2} \gamma_i \ln\left(\frac{\gamma_i}{\varepsilon c_i}\right) \leq \varepsilon \sum_{i \in I^2} c_i \quad \text{for all } \gamma_i \geq 0.$$

Thus,

$$(13) \quad v(\delta, \gamma, \varepsilon) \leq v(\delta) + \varepsilon \sum_{i \in I^2} c_i \leq f(z) + \varepsilon \sum_{i \in I^2} c_i$$

From (13) we conclude

$$(14) \quad \sup v(\delta, \gamma, \epsilon) \leq \sup v(\delta) + \epsilon \sum_{i \in I_2} c_i \leq \inf f(z) + \epsilon \sum_{i \in I_2} c_i$$

But by the theory of [1],

$$\sup v(\delta, \gamma, \epsilon) = \max v(\delta, \gamma, \epsilon) = \inf g(z, \epsilon) \geq \inf f(z) \text{ by (10.1). Hence}$$

$$(15) \quad \inf f(z) \leq \max v(\delta, \gamma, \epsilon) \leq \sup v(\delta) + \epsilon \sum_{i \in I_2} c_i \leq \inf f(z) + \epsilon \sum_{i \in I_2} c_i$$

Thus letting $\epsilon \rightarrow 0$, we have

$$(16) \quad \inf f(z) \leq \lim_{\epsilon \rightarrow 0} \max v(\delta, \gamma, \epsilon) \leq \sup v(\delta) \leq \inf f(z)$$

and

$$(17) \quad \lim_{\epsilon \rightarrow 0} \max v(\delta, \gamma, \epsilon) = \sup v(\delta) = \inf f(z)$$

Q.E.D.

REFERENCES

1. Charnes, A., Cooper, W.W., and Seiford, L., "Extremal Principles and Optimization Dualities for Khincin-Kullback-Leibler Estimation," Mathematische Operationsforschung und Statistik, Vol. 9, No. 1, 1978, p. 21-29.
2. Charnes, A., and Ben-Tal, A., "A Dual Optimization Framework for Some Problems of Information Theory and Statistics," Journal of Problems of Control and Information Theory, Vol. 8, (5-6), p. 387-401, 1979.

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER CCS 414	2. GOVT ACCESSION NO. AD-A111 326	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Khincin-Kullback-Leibler Estimation With Inequality Constraints		5. TYPE OF REPORT & PERIOD COVERED
7. AUTHOR(s) A. Charnes, W.W. Cooper, J. Tyssedal		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Center for Cybernetic Studies, UT Austin Austin, Texas 78712		8. CONTRACT OR GRANT NUMBER(s) N00014-75-C-0569
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research (Code 434) Washington, D.C.		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE November 1981
		13. NUMBER OF PAGES 9
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) This document has been approved for public release and sale; its distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Khincin-Kullback-Leibler Estimation, Inferential Distribution, Minimum Discrimination Information, Inequality Constrained Distributions		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) In this paper we extend our grasp of statistical theory, duality theory, and computational convenience to the general linear inequality situation in K^2L (or minimum discrimination information) estimation by exhibiting it as the limit of a simple one parameter sequence of equality problems. Only the finite discrete distribution case is treated here.		

DD FORM 1473
1 JAN 73

EDITION OF 1 NOV 68 IS OBSOLETE
S/N 0102-014-6601

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

ATE
LMED
-8